

Hydrogenation of a Rhodium Peroxido Complex by Formate Derivatives: Mechanistic Studies and the Catalytic Formation of H₂O₂ from O₂**

Gregor Meier and Thomas Braun*

An interesting task in oxygenation chemistry is the transition-metal-mediated hydrogenation of dioxygen.^[1] One desired product is hydrogen peroxide,^[2] which is an economically important industrial product for the manufacture of a wide range of everyday items.^[2a] At present, H₂O₂ is predominantly produced by the anthraquinone process, which involves a catalytic hydrogenation of 2-alkylanthraquinone derivatives and a subsequent oxidation of the resulting 2-alkylanthrahydroquinones with O₂.^[2a] However, the development of an alternative homogeneously catalyzed process to access H₂O₂ from O₂ can be considered as a demanding challenge.^[2]

Hence, a variety of catalytic investigations have been carried out to develop new reaction routes to access H₂O₂ from dioxygen with appropriate reducing agents.^[2–6] For example, it was reported that the hydrogenation of dioxygen gives hydrogen peroxide when using palladium complexes and alcohols as reductants.^[4] Note that copper complexes as functional models for galactose oxidase were also applied to catalyze the oxidation of alcohols with O₂, yielding the corresponding aldehydes or ketones, as well as H₂O₂.^[7] Moreover, palladium-catalyzed transformations of mixtures of CO, O₂, and H₂O provide CO₂ and H₂O₂ in the presence of an excess of a Brønsted acid.^[2b,5,6] A combination of [Pd(OAc)₂] and triphenylphosphine or triarylarsine was originally employed.^[5] The conversions are hampered by the vulnerability of the ligands towards oxygenation. However, ligand oxidation is avoided by the application of palladium complexes which bear phenanthroline ligands, and H₂O₂ is catalytically accessible under comparatively harsh reaction conditions (70 atm, O₂/CO 64:6, 70°C).^[6] Mechanistic steps might include dioxygen activation to yield a transition metal-peroxido complex,^[8] followed by protonation of the peroxido ligand to generate H₂O₂.^[2,9,10] However, a second possibility consists of the insertion of O₂ into a palladium–hydrogen bond.^[2b,9,11] Both reaction pathways involve the formation of hydroperoxido complexes as intermediates.^[10–12]


In the case of rhodium, we found in model reactions that the protonation of the peroxido complex *trans*-[Rh(O₂)(4-

C₅F₄N)(CNtBu)(PEt₃)₂] (**1a**) with HCl initially led to the formation of *trans*-[Rh(OOH)(Cl)(4-C₅F₄N)(CNtBu)(PEt₃)₂] and subsequently to H₂O₂ and the rhodium(III) complex *cis*-*trans*-[Rh(Cl)₂(4-C₅F₄N)(CNtBu)(PEt₃)₂].^[10] Complex **1a** was prepared from dioxygen and *trans*-[Rh(4-C₅F₄N)(CNtBu)-(PEt₃)₂] (**2**).

Herein, we report on the unique reactivity of **1a** towards dihydrogen sources to yield hydrogen peroxide and **2**. The metal-bound oxygen can be reduced either by the Hantzsch ethyl ester diethyl-1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylate (**3**), formic acid or ammonium formate. In a catalytic experiment, H₂O₂ was generated at ambient temperature and atmospheric pressure from dioxygen and ammonium formate.

Reaction of the peroxido complex *trans*-[Rh(O₂)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**1a**) with the Hantzsch ethyl ester **3** at 50°C led to the formation of the rhodium(I) compound *trans*-[Rh(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**2**),^[13] diethyl 2,6-dimethylpyridine-3,5-dicarboxylate (**4**) and H₂O₂ within eight hours (Scheme 1). Considerable amounts (10%) of OPEt₃ could be detected, because of the slow degradation of rhodium(I) species **2** by reaction with H₂O₂. However, in the presence of one equivalent of PPh₃, no OPEt₃ was formed and PPh₃ was instead monooxygenated to give OPPh₃.^[14]

Alternatively, the treatment of **1a** with a solution of HCOOH in THF at –70°C afforded H₂O₂, CO₂, and



Scheme 1. Formation of H₂O₂ by hydrogenation of **1a**.

[*] G. Meier, Prof. Dr. T. Braun


Humboldt-Universität zu Berlin, Department of Chemistry
Brook-Taylor-Strasse 2, 12489 Berlin (Germany)
E-mail: thomas.braun@chemie.hu-berlin.de

[**] We would like to acknowledge the Deutsche Forschungsgemeinschaft for financial support. We are grateful to S. Mebs and R. Herrmann for the crystallography.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201207073>.

rhodium(I) complex **2**, as well as the rhodium(III) carbonato complex *trans*-[Rh(κ^2 -OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**5a**; Scheme 1). The H₂O₂ can be separated as an aqueous solution by trap to trap vacuum distillation of the volatiles followed by treatment of the distillate with water. Note that in an experiment at room temperature small amounts of 2,3,5,6-tetrafluoropyridine were produced, owing to the protonation of the tetrafluoropyridyl ligand instead of the peroxydo entity.^[13,15] However, at low-temperature 2,3,5,6-tetrafluoropyridine was not formed.

Low-temperature NMR and Raman experiments at -70°C revealed the formation of the hydroperoxido formato complex *trans*-[Rh(OOH){OC(O)H}(4-C₅F₄N)(CNtBu)-(PEt₃)₂] (**6a**) as the initial product (Scheme 2). The ¹⁸O




Scheme 2. Reactivity of **1a** towards formic acid.

isotopologue *trans*-[Rh(¹⁸O¹⁸OH){OC(O)H}(4-C₅F₄N)-(CNtBu)(PEt₃)₂] (**6b**) is available in a similar fashion, starting from *trans*-[Rh(¹⁸O₂)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**1b**).^[10] In addition to the signals for **6a**, small amounts of **1a** were still detectable in the ³¹P{¹H} NMR spectrum. Further monitoring of the reaction by NMR spectroscopy at low temperature revealed that **1a** was converted into an additional intermediate species, which was identified as the peroxycarbonato compound *trans*-[Rh(κ^2 -OOC(O)O)(4-C₅F₄N)(CNtBu)-(PEt₃)₂] (**7a**; Scheme 2). We presume that initially CO₂ is generated, which reacts with **1a** to give **7a**. However, with increasing temperature, complexes **2** and **5a** were the sole rhodium species that could be identified.

The presence of the rhodium(III) complex **6a** is evidenced by a rhodium–phosphorus coupling constant of 86.3 Hz for the signal at $\delta = 22.0$ ppm in the ³¹P{¹H} NMR spectrum, and by an absorption band at 2210 cm^{-1} in the Raman spectrum, which can be assigned to the stretching vibration of the CNtBu ligand.^[10,13,16] The ¹⁹F NMR spectrum of **6a** displays four signals for the tetrafluoropyridyl ligand, which indicates a hindered rotation about the rhodium–carbon bond.^[10,13,16] A broad signal in the ¹H NMR spectrum at $\delta = 8.38$ ppm can be attributed to the hydroperoxido ligand,^[10,12,17] and a resonance

at $\delta = 8.15$ ppm is assigned to the formato ligand.^[18] The latter signal correlates with a signal for the {OC(O)H} ligand at $\delta = 167.5$ ppm in the ¹³C domain of a ¹H,¹³C-HMQC NMR spectrum.^[18] Furthermore, the Raman spectrum of **6a** shows an absorption band at 1624 cm^{-1} for the formato ligand. Another band at 859 cm^{-1} shifts to 815 cm^{-1} for the ¹⁸O-labeled isotopologue **6b** ($\Delta\nu = 44\text{ cm}^{-1}$) and can be assigned to the OOH stretching vibration.^[12a,19]

There is some precedence for late transition metal compounds reacting with O₂ and subsequently CO₂ to yield peroxycarbonato complexes.^[20–22] To get a deeper understanding of the formation and reactivity of complexes **5a** and **7a**, the latter was synthesized in an independent synthesis. Thus, a solution of *trans*-[Rh(O₂)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**1a**) in benzene was treated with CO₂ to give the peroxycarbonato compound *trans*-[Rh(κ^2 -OOC(O)O)(4-C₅F₄N)-(CNtBu)(PEt₃)₂] (**7a**; Scheme 3). The compounds *trans*-[Rh(κ^2 -¹⁸O¹⁸OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**7b**) and



Scheme 3. Formation and reactivity of the peroxycarbonato complex **7a** and carbonato complex **5a**.

trans-[Rh(κ^2 -¹⁷O¹⁷OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**7c**) were available in a similar fashion from the ¹⁸O- and ¹⁷O-labeled isotopomers **1b** and **1c**, respectively. Treatment of the ¹⁷O-labeled peroxydo complex **1c** with ¹³CO₂ afforded the ¹³C-enriched isotopologue *trans*-[Rh(κ^2 -¹⁷O¹⁷O¹³C(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**7d**).

The carbonato complex *trans*-[Rh(κ^2 -OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**5a**) and its ¹⁸O isotopologue *trans*-[Rh(κ^2 -¹⁸OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**5b**) were also synthesized by an independent reaction pathway. The treatment of **7a** or **7b** with PPh₃ led to a monooxygenation of the phosphine to form OPPh₃ and the carbonato complexes *trans*-[Rh(κ^2 -OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**5a**) and *trans*-[Rh(κ^2 -¹⁸OC(O)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**5b**), respectively (Scheme 3). Remarkably, complex **7a** also reacts in an unprecedented reaction with aqueous H₂O₂ to yield **5a**, water, and O₂ (Scheme 3). Thus overall, the reaction of **1a** with

formic acid initially gives H_2O_2 and CO_2 , as well as **2** (Scheme 2). Carbon dioxide reacts subsequently with **1a** to generate the peroxycarbonato compound **7a**, which yielded the carbonate **5a**, H_2O , and O_2 by reaction with H_2O_2 (Scheme 3). Above all, it is intriguing that the rhodium(III) compounds **5a** and **7a** also both react with HCOOH to give the rhodium(I) complex **2** (Scheme 3).

A signal in the $^{31}\text{P}\{\text{H}\}$ NMR spectrum for **5a** at $\delta = 23.0$ ppm exhibits a rhodium–phosphorus coupling constant of 83.4 Hz. An absorption band at 2194 cm^{-1} in the IR spectrum (ATR) corresponds to the stretching vibration of the CNtBu ligand at a rhodium(III) complex.^[10,13,16] The absorption bands at 673 cm^{-1} for **5a**, as well as 674 cm^{-1} , 662 cm^{-1} , and 649 cm^{-1} for **5b** can tentatively be assigned to in-plane deformation vibrations of the $\text{Rh}\{\kappa^2-\text{OC(O)O}\}$ moiety.^[23] The structure of **5a** in the solid state was determined by X-ray crystallography (Figure 1).^[24] The Rh1–O1 and Rh1–O2 distances (2.0625(17) Å and




Figure 1. Crystal structure of **5a** (ORTEP, ellipsoids set at 50% probability). Selected distances [Å] and angles [°]: C1–Rh1 2.060(3), C18–Rh1 1.926(3), O1–Rh1 2.0625(17), O2–Rh1 2.0856(17), C23–O1 1.322(3), C23–O2 1.320(3), C23–O3 1.222(3); N2–C18–Rh1 175.3(2), C18–Rh1–C1 91.72(10), C18–Rh1–O1 165.70(9), C1–Rh1–O1 102.54(9), O1–Rh1–O2 63.75(7), O2–C23–O1 112.0(2).

2.0856(17) Å) are equivalent to those found in *trans*-[$\text{Rh}\{\kappa^2-\text{OC(O)O}\}(\text{S}_2\text{CNMe}_2)(\text{PPh}_3)_2$]^[20a] (2.071(8) Å and 2.083(8) Å); the O–Rh–O angle of 63.3(3)° in the latter is similar to the O1–Rh1–O2 angle of 63.75(7)° in **5a**.

The presence of rhodium(III) complex **7a** is again evidenced by a rhodium–phosphorus coupling constant of 85.5 Hz for the signal at $\delta = 22.9$ ppm in the $^{31}\text{P}\{\text{H}\}$ NMR spectrum and by an absorption band at 2191 cm^{-1} in the IR spectrum (ATR), which can be assigned to the stretching vibration of the CNtBu ligand.^[10,13,16] The ^{17}O NMR spectrum of **7c** displays two signals (Figure 2a). A signal at $\delta = 38.7$ ppm with a rhodium–oxygen coupling constant of 150.0 Hz can be attributed to the metal bound oxygen atom of the ^{17}O – ^{17}O moiety in the peroxycarbonato ligand. A




Figure 2. a) ^{17}O NMR spectrum of *trans*-[$\text{Rh}\{\kappa^2-\text{¹⁷O}^{¹⁷}\text{OC(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$] (**7c**); b) Part of the ^{17}O NMR spectrum of *trans*-[$\text{Rh}\{\kappa^2-\text{¹⁷O}^{¹⁷}\text{O}^{¹³}\text{C(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$] (**7d**).

resonance at $\delta = 76.2$ ppm is assigned to the carbon-bound oxygen atom of the ^{17}O – ^{17}O moiety. The ^{17}O NMR spectrum of the ^{13}C isotopologue **7d** reveals a carbon–oxygen coupling constant of 15.3 Hz for the latter resonance (Figure 2b). The IR spectrum of **7a** shows an absorption band at 969 cm^{-1} for the $\text{Rh}\{\kappa^2-\text{OOC(O)O}\}$ moiety.^[20b] This band shifts to 959 cm^{-1} for the ^{17}O -labeled isotopologue **7c** ($\Delta\nu = 10\text{ cm}^{-1}$) and to 948 cm^{-1} for the ^{18}O -labeled isotopologue **7b** ($\Delta\nu = 21\text{ cm}^{-1}$), respectively (Table 1).

Table 1: Selected IR spectroscopic data (ATR) for the peroxycarbonato complexes **7a**–**7d**.

Entry	Complex	$\tilde{\nu}$ [cm ⁻¹] ^[a]	$\Delta\nu$ [cm ⁻¹] ^[b]	$\tilde{\nu}$ [cm ⁻¹] ^[b]	$\Delta\nu$ [cm ⁻¹] ^[b]
1	7a	969	–	585; 543	–
2	7b	948	21 ^[c]	566; 519	19; 24 ^[c]
3	7c	959	10 ^[c]	575; 531	10; 12 ^[c]
4	7d	952	7 ^[d]	575; 531	0; 0 ^[d]

[a] Vibrational frequencies of the $\text{Rh}\{\kappa^2-\text{OOC(O)O}\}$ moiety. [b] Vibrational frequencies of the RhO moiety. [c] Isotopic shifts compared to **7a**. [d] Isotopic shifts compared to **7c**. **7a**: *trans*-[$\text{Rh}\{\kappa^2-\text{OOC(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$], **7b**: *trans*-[$\text{Rh}\{\kappa^2-\text{¹⁷O}^{¹⁸}\text{O}^{¹³}\text{C(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$], **7c**: *trans*-[$\text{Rh}\{\kappa^2-\text{¹⁷O}^{¹⁷}\text{OC(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$], **7d**: *trans*-[$\text{Rh}\{\kappa^2-\text{¹⁷O}^{¹⁷}\text{O}^{¹³}\text{C(O)O}\}(4-\text{C}_5\text{F}_4\text{N})(\text{CNtBu})(\text{PEt}_3)_2$].

These frequency shifts are too small to be associated with isolated $\nu(\text{O}-\text{O})$ modes in the peroxycarbonato ligands of **7a**, **7b**, and **7c**, respectively.^[20b] Moreover, the IR spectrum of **7d** shows an absorption band at 952 cm^{-1} . Thus, the absorption band at 959 cm^{-1} for the ^{17}O -labeled isotopologue **7c** displays an isotopic shift of $\Delta\nu = 7\text{ cm}^{-1}$ upon ^{13}C labeling. Absorption bands at 585 cm^{-1} and 543 cm^{-1} for **7a** shift to 575 cm^{-1} ($\Delta\nu = 10\text{ cm}^{-1}$) and 531 cm^{-1} ($\Delta\nu = 12\text{ cm}^{-1}$) for **7c** and to 566 cm^{-1} ($\Delta\nu = 19\text{ cm}^{-1}$) and 519 cm^{-1} ($\Delta\nu = 24\text{ cm}^{-1}$) for **7b**. These bands do not show any isotopic shift upon ^{13}C labeling and can tentatively be associated with vibrations which include the RhO moiety.^[20b] Based on the ^{17}O NMR and IR data, we suggest that CO_2 reacts with rhodium peroxydo complexes by insertion into a metal–oxygen^[22] bond rather than by insertion

into the O–O bond, as was previously suggested.^[20b,c] A comparable reactivity was found in the reaction of boronic acids with **1a**, in which *trans*-[Rh(κ^2 -OOB(C₆R₅)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (R = H, F) are formed by cleavage of the Rh–O bond of the peroxydo moiety in **1a**.^[16a]

The molecular structure of **7a**·H₂O in the solid state was determined by X-ray crystallography (Figure 3).^[24] The mol-





Figure 3. Crystal structure of **7a**·H₂O (ORTEP, ellipsoids set at 50% probability). Complex **7a** crystallizes with an additional water molecule, which is hydrogen bonded to the peroxycarbonato ligand. Selected distances [Å] and angles [°]: Rh1–C2 1.943(4), Rh1–C7 2.085(4), Rh1–O1 2.040(3), Rh1–O3 2.081(3), O1–O2 1.469(4), C1–O2 1.354(6), C1–O3 1.255(5), C1–O4 1.238(5); N1–C2–Rh1 175.1(4), C2–Rh1–C7 90.80(16), C2–Rh1–O1 171.48(14), O1–Rh1–C7 97.72(14), C2–Rh1–O3 92.85(14), O1–Rh1–O3 78.65(12).

ecule exhibits a distorted octahedral geometry with the two phosphines in a mutually *trans* orientation. The O1–Rh1–O3 angle of 78.65(12)° is comparable to the corresponding angle in the peroxycarbonatocomplex [Rh(κ^2 -OOC(O)O)(4-MeC₆H₄){tBuP(CH₂CH₂CH₂-PPh₂)₂}].^[21a] The O–O distance of 1.49(2) Å in the latter,^[21a] as well as the O–O distance of 1.4974(14) Å in *trans*-[Rh(κ^2 -OOB(OH)O)(4-C₅F₄N)(CNtBu)(PEt₃)₂]^[16a] are comparable to the O1–O2 bond length of 1.469(4) Å in **7a**. The O–O distances in other cyclic peroxycarbonatocomplexes are also in a similar range, as found for **7a**.^[21]

In the next step we applied ammonium formate as dihydrogen source and as a mild alternative to formic acid.^[25,26] Notably, reaction with **1a** afforded H₂O₂, CO₂, NH₃, and the rhodium complexes **2** and **5a** at ambient temperature (Scheme 1). We then tested the reactivity of dioxygen towards ammonium formate in the presence of substoichiometric amounts of the peroxydo complex **1a** (Scheme 4). Indeed, dioxygen and ammonium formate can be transformed into H₂O₂, NH₃, and CO₂ in the presence of **1a** (20%) at ambient temperature and atmospheric pressure. The generated hydrogen peroxide can be utilized in oxygenation reactions, such as the *in situ* oxygenation of PPh₃.^[14]



Scheme 4. Formation of H₂O₂ with substoichiometric amounts of **1a**. DABCO = 1,4-diazabicyclo[2.2.2]octane.

Furthermore, H₂O₂ was trapped as the adduct (H₂O₂)₂·DABCO (DABCO = 1,4-diazabicyclo[2.2.2]octane; Scheme 4). Based on **1a**, 3.6 equiv of H₂O₂ were formed.^[27]

In conclusion, we have demonstrated a variety of hydrogenation reactions of the peroxydo ligand in *trans*-[Rh(O₂)(4-C₅F₄N)(CNtBu)(PEt₃)₂] (**1a**), which can be prepared from dioxygen and **2**, to yield H₂O₂ and the rhodium(I) compound **2**. To the best of our knowledge, the hydrogenation of dioxygen with formic acid or ammonium formate in the homogeneous phase mediated by a transition metal complex is unprecedented.^[28] The reduction of dioxygen at ambient temperature and atmospheric pressure with ammonium formate yielding H₂O₂ can be mediated by substoichiometric amounts of **1a**.

Received: August 31, 2012

Published online: November 4, 2012

Keywords: hydrogenation · hydrogen peroxide · oxidation · peroxydo ligands · rhodium

- [1] See, for example: a) *Topics in Organometallic Chemistry* (Eds.: C. Limberg, F. Meyer), Springer, Berlin, 2007; b) *Modern Oxidation Methods* (Ed.: J.-E. Bäckvall), Wiley-VCH, Weinheim, 2004; c) C. Limberg, *Angew. Chem.* **2003**, *115*, 6112–6136; *Angew. Chem. Int. Ed.* **2003**, *42*, 5932–5954; d) J.-M. Brégeault, *Dalton Trans.* **2003**, 3289–3302; e) E. C. Niederhoffer, J. H. Timmons, A. E. Martell, *Chem. Rev.* **1984**, *84*, 137–203; f) R. A. Sheldon, J. K. Kochi, *Metal-Catalyzed Oxidations of Organic Compounds*, Academic Press, New York, 1981; g) S. S. Stahl, *Science* **2005**, *309*, 1824–1826; h) L. Boisvert, K. I. Goldberg, *Acc. Chem. Res.* **2012**, *45*, 899–910; i) A. N. Campbell, S. S. Stahl, *Acc. Chem. Res.* **2012**, *45*, 851–863; j) A. N. Vedernikov, *Acc. Chem. Res.* **2012**, *45*, 803–813.
- [2] See, for example: a) J. M. Campos-Martin, G. Blanco-Brieva, J. L. G. Fierro, *Angew. Chem.* **2006**, *118*, 7116–7139; *Angew. Chem. Int. Ed.* **2006**, *45*, 6962–6984; b) W. R. Thiel, *Angew. Chem.* **1999**, *111*, 3349–3351; *Angew. Chem. Int. Ed.* **1999**, *38*, 3157–3158; c) S. S. Stahl, *Angew. Chem.* **2004**, *116*, 3480–3501; *Angew. Chem. Int. Ed.* **2004**, *43*, 3400–3420; d) E. G. Chepaikin, *Russ. Chem. Rev.* **2011**, *80*, 363–396.
- [3] See, for example: a) J. K. Edwards, G. J. Hutchings, *Angew. Chem.* **2008**, *120*, 9332–9338; *Angew. Chem. Int. Ed.* **2008**, *47*, 9192–9198; b) P. Landon, P. J. Collier, A. F. Carley, D. Chadwick, A. J. Papworth, A. Burrows, C. J. Kiely, G. J. Hutchings, *Phys. Chem. Chem. Phys.* **2003**, *5*, 1917–1923; c) J. K. Edwards, B. Solsona, P. Landon, A. F. Carley, A. Herzing, M. Watanabe, C. J. Kiely, G. J. Hutchings, *J. Mater. Chem.* **2005**, *15*, 4595–4600; d) R. Burch, P. R. Ellis, *Appl. Catal. B* **2003**, *42*, 203–211; e) J. K. Edwards, B. E. Solsona, P. Landon, A. F. Carley, A. Herzing, C. J.

- Kiely, G. J. Hutchings, *J. Catal.* **2005**, 236, 69–79; f) S. Park, T. J. Kim, Y.-M. Chung, S.-H. Oh, I. K. Song, *Korean J. Chem. Eng.* **2011**, 28, 1359–1363; g) Y. Zuo, J. Holgné, *Environ. Sci. Technol.* **1992**, 26, 1014–1022; h) B. Su, R. P. Nia, F. Li, M. Hojeij, M. Prudent, C. Corminboeuf, Z. Samec, H. H. Girault, *Angew. Chem.* **2008**, 120, 4753–4756; *Angew. Chem. Int. Ed.* **2008**, 47, 4675–4678; i) S. Chowdhury, F. Himo, N. Russo, E. Sicilia, *J. Am. Chem. Soc.* **2010**, 132, 4178–4190.
- [4] R. Bortolo, D. Bianchi, R. D'Aloisio, C. Querci, M. Ricci, *J. Mol. Catal. A* **2000**, 153, 25–29.
- [5] a) V. N. Zudin, V. A. Likholobov, Y. I. Ermakov, *Kinet. Katal.* **1977**, 18, 921–926; b) V. N. Zudin, V. A. Likholobov, Y. I. Ermakov, *Kinet. Katal.* **1979**, 20, 1599–1600.
- [6] a) D. Bianchi, R. Bortolo, R. D'Aloisio, M. Ricci, *Angew. Chem.* **1999**, 111, 734–736; *Angew. Chem. Int. Ed.* **1999**, 38, 706–708; b) D. Bianchi, R. Bortolo, R. D'Aloisio, M. Ricci, *J. Mol. Catal. A* **1999**, 150, 87–94; c) D. Bianchi, R. Bortolo, R. D'Aloisio, C. Querci, M. Ricci, *Stud. Surf. Sci. Catal.* **1999**, 126, 481–484; d) C. Querci, R. D'Aloisio, R. Bortolo, M. Ricci, D. Bianchi, *J. Mol. Catal. A* **2001**, 176, 95–100.
- [7] a) C. T. Lyons, T. D. P. Stack, *Coord. Chem. Rev.*, DOI: 10.1016/j.ccr.2012.06.003; b) B. A. Jazdzewski, W. B. Tolman, *Coord. Chem. Rev.* **2000**, 200–202, 633–685; c) P. Chaudhuri, M. Hess, T. Weyhermüller, K. Wieghardt, *Angew. Chem.* **1999**, 111, 1165–1168; *Angew. Chem. Int. Ed.* **1999**, 38, 1095–1098; d) P. Chaudhuri, M. Hess, U. Flörke, K. Wieghardt, *Angew. Chem.* **1998**, 110, 2340–2343; *Angew. Chem. Int. Ed.* **1998**, 37, 2217–2220; e) P. Chaudhuri, M. Hess, J. Müller, K. Hildenbrandt, E. Bill, T. Weyhermüller, K. Wieghardt, *J. Am. Chem. Soc.* **1999**, 121, 9599–9610; f) T. K. Paine, T. Weyhermüller, K. Wieghardt, P. Chaudhuri, *Dalton Trans.* **2004**, 2092–2101; g) Y. Wang, T. D. P. Stack, *J. Am. Chem. Soc.* **1996**, 118, 13097–13098; h) B. A. Jazdzewski, V. G. Young, Jr., W. B. Tolman, *Chem. Commun.* **1998**, 2521–2522.
- [8] For some recently reported rhodium peroxydo complexes, see: a) T. S. Teets, D. G. Nocera, *Inorg. Chem.* **2012**, 51, 7192–7201; b) D. Inoki, T. Matsumoto, H. Hayashi, K. Takashita, H. Nakai, S. Ogo, *Dalton Trans.* **2012**, 41, 4328–4334; c) G. Meier, T. Braun, *Angew. Chem.* **2011**, 123, 3338–3342; *Angew. Chem. Int. Ed.* **2011**, 50, 3280–3284; d) A. Penner, T. Braun, *Eur. J. Inorg. Chem.* **2011**, 2579–2587; e) A. Penner, T. Schröder, T. Braun, B. Ziemer, *Eur. J. Inorg. Chem.* **2009**, 4464–4470; f) J. Vicente, J. Gil-Rubio, J. Guerrero-Leal, D. Bautista, *Dalton Trans.* **2009**, 3854–3866; g) A. Y. Verat, H. Fan, M. Pink, Y.-S. Chen, K. G. Caulton, *Chem. Eur. J.* **2008**, 14, 7680–7686; h) X.-Y. Yu, B. O. Patrick, B. R. James, *Organometallics* **2006**, 25, 4870–4877; i) J. R. Bleeke, M. Shokeen, E. S. Wise, N. P. Rath, *Organometallics* **2006**, 25, 2486–2500; j) B. de Bruin, T. P. J. Peters, J. B. M. Wilting, S. Thewissen, J. M. M. Smits, A. W. Gal, *Eur. J. Inorg. Chem.* **2002**, 2671–2680; k) D. G. Ho, R. Ismail, N. Franco, R. Gao, E. P. Leverich, I. Tsypa, N. N. Ho, R. Bau, M. Selke, *Chem. Commun.* **2002**, 570–571; l) L. Carlton, L. V. Mokoena, M. A. Fernandes, *Inorg. Chem.* **2008**, 47, 8696–8703; m) O. V. Zenkina, E. C. Keske, R. Wang, C. M. Crudden, *Angew. Chem.* **2011**, 123, 8250–8254; *Angew. Chem. Int. Ed.* **2011**, 50, 8100–8104.
- [9] a) M. M. Konnick, S. S. Stahl, *J. Am. Chem. Soc.* **2008**, 130, 5753–5762; b) B. V. Popp, S. S. Stahl, *J. Am. Chem. Soc.* **2007**, 129, 4410–4422; c) M. M. Konnick, B. A. Gandhi, I. A. Guzei, S. S. Stahl, *Angew. Chem.* **2006**, 118, 2970–2973; *Angew. Chem. Int. Ed.* **2006**, 45, 2904–2907.
- [10] M. Ahijado, T. Braun, D. Noveski, N. Kocher, B. Neumann, D. Stalke, H.-G. Stammmer, *Angew. Chem.* **2005**, 117, 7107–7111; *Angew. Chem. Int. Ed.* **2005**, 44, 6947–6951.
- [11] a) T. Hosokawa, S.-I. Murahashi, *Acc. Chem. Res.* **1990**, 23, 49–54; b) T. Nishimura, T. Onoue, K. Ohe, S. Uemura, *J. Org. Chem.* **1999**, 64, 6750–6755; c) D. D. Wick, K. I. Goldberg, *J. Am. Chem. Soc.* **1999**, 121, 11900–11901; d) M. C. Denney, N. A. Smythe, K. L. Cetto, R. A. Kemp, K. I. Goldberg, *J. Am. Chem. Soc.* **2006**, 128, 2508–2509; e) J. M. Keith, R. P. Muller, R. A. Kemp, K. I. Goldberg, W. A. Goddard III, J. Oxgaard, *Inorg. Chem.* **2006**, 45, 9631–9633; f) T. Hosokawa, T. Nakahira, M. Takano, S. Murahashi, *J. Mol. Catal.* **1992**, 74, 489–498; g) T. Nishimura, N. Kakiuchi, T. Onoue, K. Ohe, S. Uemura, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1915–1918; h) M. M. Konnick, N. Decharin, B. V. Popp, S. S. Stahl, *Chem. Sci.* **2011**, 2, 326–330; i) K. M. Gligorich, M. S. Sigman, *Angew. Chem.* **2006**, 118, 6764–6767; *Angew. Chem. Int. Ed.* **2006**, 45, 6612–6615; see also, Ref. [9b].
- [12] a) D. Carmona, M. P. Lamata, J. Ferrer, J. Modrego, M. Perales, F. J. Lahoz, R. Atencio, L. A. Oro, *J. Chem. Soc. Chem. Commun.* **1994**, 575–576; b) C. Tejel, M. A. Ciriano, S. Jiménez, V. Passarelli, J. A. López, *Angew. Chem.* **2008**, 120, 2123–2126; *Angew. Chem. Int. Ed.* **2008**, 47, 2093–2096; c) M. M. Konnick, I. A. Guzei, S. S. Stahl, *J. Am. Chem. Soc.* **2004**, 126, 10212–10213; d) Y. Takahashi, M. Hashimoto, S. Hikichi, M. Akita, Y. Moro-oka, *Angew. Chem.* **1999**, 111, 3259–3262; *Angew. Chem. Int. Ed.* **1999**, 38, 3074–3077; e) V. V. Rostovtsev, L. M. Henling, J. A. Labinger, J. E. Bercaw, *Inorg. Chem.* **2002**, 41, 3608–3619; f) M. Krom, T. P. J. Peters, R. G. E. Coumans, T. J. J. Sciarone, J. Hoogboom, S. I. ter Beek, P. P. J. Schlebos, J. M. M. Smits, R. de Gelder, A. W. Gal, *Eur. J. Inorg. Chem.* **2003**, 1072–1087.
- [13] D. Noveski, T. Braun, B. Neumann, A. Stammmer, H.-G. Stammmer, *Dalton Trans.* **2004**, 4106–4119.
- [14] a) C. W. Cheung, K. S. Chan, *Organometallics* **2011**, 30, 4999–5009; b) A. Cervilla, F. Pérez-Plá, E. Llopis, M. Piles, *Dalton Trans.* **2004**, 1461–1465; c) M. M. Abu-Omar, J. H. Espenson, *J. Am. Chem. Soc.* **1995**, 117, 272–280.
- [15] a) T. Braun, D. Noveski, M. Ahijado, F. Wehmeier, *Dalton Trans.* **2007**, 3820–3825; b) A referee pointed out that complex **2** might react with HCOOH to give an oxidative addition product and subsequently CO by dehydration. We have no spectroscopic indication (NMR, IR, mass spectrometry) for these reactions; see also: W. I. Dzik, J. M. M. Smits, J. N. H. Reek, B. de Bruin, *Organometallics* **2009**, 28, 1631–1643.
- [16] a) M. Ahijado, T. Braun, *Angew. Chem.* **2008**, 120, 2996–3000; *Angew. Chem. Int. Ed.* **2008**, 47, 2954–2958; b) M. Ahijado Salomon, T. Braun, A. Penner, *Angew. Chem.* **2008**, 120, 8999–9003; *Angew. Chem. Int. Ed.* **2008**, 47, 8867–8871.
- [17] a) I. A. Guzei, A. Bakac, *Inorg. Chem.* **2001**, 40, 2390–2393; b) T. Miyaji, M. Kujime, S. Hikichi, Y. Moro-oka, M. Akita, *Inorg. Chem.* **2002**, 41, 5286–5295; see also, Ref. [11c].
- [18] a) Y. S. Varshavskii, T. G. Cherkasova, I. S. Podkorytov, A. A. Korlyukov, V. N. Khrustalev, A. B. Nikol'skii, *Russ. J. Coord. Chem.* **2005**, 31, 121–131; b) W. Keim, J. Becker, *J. Organomet. Chem.* **1989**, 372, 447–451; c) V. V. Grushin, V. F. Kuznetsov, C. Bensimon, H. Alper, *Organometallics* **1995**, 14, 3927–3932; d) F. P. Pruchnik, A. Jutarska, Z. Ciunik, M. Pruchnik, *Inorg. Chim. Acta* **2003**, 350, 609–616.
- [19] a) H. L. Roberts, W. R. Symes, *J. Chem. Soc. A* **1968**, 6, 1450–1453; b) H. Suzuki, S. Matsuura, Y. Moro-oka, T. Ikawa, *J. Organomet. Chem.* **1985**, 286, 247–258.
- [20] See, for example: a) Y. Wakatsuki, M. Maniwa, H. Yamazaki, *Inorg. Chem.* **1990**, 29, 4204–4208; b) M. Aresta, I. Tommasi, E. Quaranta, C. Fragale, J. Mascetti, M. Tranquille, F. Galan, M. Fouassier, *Inorg. Chem.* **1996**, 35, 4254–4260; c) M. Aresta, A. Dibenedetto, I. Tommasi, *Eur. J. Inorg. Chem.* **2001**, 1801–1806; d) M. G. Mason, J. A. Ibers, *J. Am. Chem. Soc.* **1982**, 104, 5153–5157; e) H. J. Lawson, J. D. Atwood, *J. Am. Chem. Soc.* **1989**, 111, 6223–6227; f) P. J. Hayward, D. M. Blake, G. Wilkinson, C. J. Nyman, *J. Am. Chem. Soc.* **1970**, 92, 5873–5878; g) G. Paiaro, L. Pandolfo, F. Morandini, S. D. Fioretto, *J. Organomet. Chem.* **1994**, 483, 147–151.

- [21] a) L. Dahlenburg, C. Prengel, *J. Organomet. Chem.* **1986**, *308*, 63–71; b) M. Yamashita, K. Goto, T. Kawashima, *J. Am. Chem. Soc.* **2005**, *127*, 7294–7295; c) A. Wada, S. Yamaguchi, K. Jitsukawa, H. Masuda, *Angew. Chem.* **2005**, *117*, 5844–5847; *Angew. Chem. Int. Ed.* **2005**, *44*, 5698–5701; d) K. Hashimoto, S. Nagatomo, S. Fucinami, H. Furutachi, S. Ogo, M. Suzuki, A. Uehara, Y. Maeda, Y. Watanabe, K. Kitagawa, *Angew. Chem.* **2002**, *114*, 1250–1253; *Angew. Chem. Int. Ed.* **2002**, *41*, 1202–1205; e) L. Dahlenburg, C. Prengel, *Organometallics* **1984**, *3*, 934–936; f) F. Hinrichs, A. Adam, *Z. Anorg. Allg. Chem.* **2011**, *637*, 426–429.
- [22] See Ref. [20g].
- [23] a) H. G. Thode, M. Shimac, C. E. Rees, K. V. Krishnamurty, *Can. J. Chem.* **1965**, *43*, 582–595; b) K. V. Krishnamurty, G. M. Harris, V. S. Sastri, *Chem. Rev.* **1970**, *70*, 171–197; c) E. Uzunova, D. Klissurski, I. Mitov, P. Stefanov, *Chem. Mater.* **1993**, *5*, 576–582; d) A. Markovits, M. García-Hernández, J. M. Ricart, F. Illas, *J. Phys. Chem. B* **1999**, *103*, 509–518; e) R. E. Hester, W. E. L. Grossman, *Inorg. Chem.* **1966**, *5*, 1308–1312; f) M. A. Andrews, G. L. Gould, W. T. Klooster, K. S. Koenig, E. J. Voss, *Inorg. Chem.* **1996**, *35*, 5478–5483; g) F. Porta, F. Ragaini, S. Cenini, *Inorg. Chim. Acta* **1990**, *173*, 229–235.
- [24] Data for X-ray structure analysis: All structures were solved by direct methods and refined with the full matrix least square methods on F^2 (SHELX-97).^[29] **5a:** $C_{23}H_{39}F_4N_2O_3P_2Rh$, $M = 632.41$, crystal dimensions $0.42 \times 0.35 \times 0.23 \text{ mm}^3$; monoclinic; $P2_1/c$; $a = 9.9319(3)$, $b = 32.7822(8)$, $c = 9.5360(3) \text{ \AA}$, $\beta = 113.617(2)^\circ$, $Z = 4$, $V = 2844.77(14) \text{ \AA}^3$, $\rho_{\text{calcd.}} = 1.477 \text{ g cm}^{-3}$; $2\theta_{\text{max}} = 54.46^\circ$, Mo $K\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$), $T = 100(2) \text{ K}$, 39170 reflections collected, 6305 were unique ($R_{\text{int.}} = 0.0707$); STOE IPDS 2T diffractometer; multiscan absorption correction (min./max. transmission 0.8728/0.7162), $\mu = 0.765 \text{ mm}^{-1}$. Final R_1 , wR_2 values on all data: 0.0408, 0.0656; R_1 , wR_2 values for 6305 reflections with $I_o > 2\sigma(I_o)$: 0.0287, 0.0644; residual electron density +0.583/−0.874 e \AA^{-3} . **7a·H₂O:** $C_{23}H_{41}F_4N_2O_3P_2Rh$, $M = 666.43$, crystal dimensions $0.13 \times 0.09 \times 0.02 \text{ mm}^3$; triclinic; $P\bar{1}$; $a = 9.5236(5)$, $b = 9.9425(5)$, $c = 17.3346(10) \text{ \AA}$, $\alpha = 78.518(4)$, $\beta = 82.396(4)$, $\gamma = 65.179(4)^\circ$, $Z = 2$, $V = 1457.75(13) \text{ \AA}^3$, $\rho_{\text{calcd.}} = 1.518 \text{ g cm}^{-3}$; $2\theta_{\text{max}} = 58.00^\circ$, Mo $K\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$), $T = 100(2) \text{ K}$, 15384 reflections collected, 6572 were unique ($R_{\text{int.}} = 0.0475$); STOE IPDS 2T diffractometer; multiscan absorption correction (min./max. transmission 0.9749/0.8534), $\mu = 0.756 \text{ mm}^{-1}$. Final R_1 , wR_2 values on all data: 0.0662, 0.1236.
- R_1 , wR_2 values for 6572 reflections with $I_o > 2\sigma(I_o)$: 0.0446, 0.1118; residual electron density +0.756/−0.998 e \AA^{-3} . CCDC 899217 (**5a**) and 899218 (**7a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [25] a) S. Ram, R. E. Ehrenhauser, *Synthesis* **1988**, 91–95; b) H. C. Lo, O. Buriez, J. B. Kerr, R. H. Fish, *Angew. Chem.* **1999**, *111*, 1524–1527; *Angew. Chem. Int. Ed.* **1999**, *38*, 1429–1432; c) D. Westerhausen, S. Herrmann, W. Hummel, E. Steckhan, *Angew. Chem.* **1992**, *104*, 1496–1498; *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 1529–1531.
- [26] For the hydrogenation of CO₂ see, for example: a) T. Schaub, R. A. Paciello, *Angew. Chem.* **2011**, *123*, 7416–7420; *Angew. Chem. Int. Ed.* **2011**, *50*, 7278–7282; b) W. Wang, S. Wang, X. Ma, J. Gong, *Chem. Soc. Rev.* **2011**, *40*, 3703–3727; c) T. Reda, C. M. Plugge, N. J. Abram, J. Hirst, *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 10654–10658; d) P. G. Jessop, T. Ikariya, R. Noyori, *Nature* **1994**, *368*, 231–233; e) A. Boddien, Felix Gärtner, C. Federsel, P. Sponholz, D. Mellmann, R. Jackstell, H. Junge, M. Beller, *Angew. Chem.* **2011**, *123*, 6535–6538; *Angew. Chem. Int. Ed.* **2011**, *50*, 6411–6414; f) A. Robertson, T. Matsumoto, S. Ogo, *Dalton Trans.* **2011**, *40*, 10304–10310; g) C. Federsel, R. Jackstell, M. Beller, *Angew. Chem.* **2010**, *122*, 6392–6395; *Angew. Chem. Int. Ed.* **2010**, *49*, 6254–6257; h) S. Enthalter, J. von Langermann, T. Schmidt, *Energy Environ. Sci.* **2010**, *3*, 1207–1217; i) S. Wesselbaum, U. Hintermair, W. Leitner, *Angew. Chem.* **2012**, *124*, 8713–8716; *Angew. Chem. Int. Ed.* **2012**, *51*, 8585–8588.
- [27] The H₂O₂ was also extracted with water, and the amount was determined by titrating with an alkaline aqueous solution of [Fe(CN)₆]^{3−} and measuring the decrease of the absorption band in the UV/Vis spectrum: a) F. Aziz, G. A. Mirza, *Talanta* **1964**, *11*, 889–892; b) J. W. Kee, Y. Y. Tan, B. H. G. Swennenhuys, A. A. Bengali, W. Y. Fan, *Organometallics* **2011**, *30*, 2154–2159.
- [28] The hydrogenation of dioxygen with formic acid as the H₂ source can be catalyzed by heterogeneous palladium catalysts, see: M. S. Yalfani, S. Contreras, F. Medina, J. Sueiras, *Chem. Commun.* **2008**, 3885–3887.
- [29] a) SHELX-97, G. M. Sheldrick, program for crystal structure refinement, University of Göttingen, **1997**; b) G. M. Sheldrick, *Acta Crystallogr. Sect. A* **2008**, *64*, 112–122.