

CONTENTS

| | |
|--|-----|
| A. TUMENDELGER, S. TOYODA, N. YOSHIDA, H. SHIOMI and R. KOUNO: Isotopocule characterization of N ₂ O dynamics during simulated wastewater treatment under oxic and anoxic conditions | 105 |
| A. AIUPPA, E. LO COCO, M. LIUZZO, G. GIUDICE, G. GIUFFRIDA and R. MORETTI: Terminal Strombolian activity at Etna's central craters during summer 2012: The most CO ₂ -rich volcanic gas ever recorded at Mount Etna | 123 |
| Z.-H. YE, P. WANG, X.-K. XIANG, A.-J. WANG, Q. YAN and Y.-K. LI: Re-Os dating and H-O isotope geochemistry of the Shimensi deposit, northern Jiangxi, China: Implication for ore genesis | 139 |
| X. SU, E. ZUO, H. LV, Q. ZHAO, P. ZHU, G. LIN and M. LIU: Fate and reactions of methane during biodegradation in an aquifer contaminated with petro- leum hydrocarbons in Northeast China | 153 |
| H. XU, D. HOU, B. CAO, X. CHEN and T. XU: Geochemical characteristics of high-maturity crude oils in the Xihu Depression, East China Sea Basin | 163 |
| R. AKHTER, N. SHIRAI and M. EBIHARA: Chemical characterization of a chromitite reference sample GPt-5 using INAA and ICP-MS | 179 |
| X. SHAO, W. YANG, W. LIANG and M. WU: Soil respiration dynamics in typical tidal flat wetlands of Hangzhou Bay, China | 187 |
| S. A. MAHMUD, S. NASEEM, M. HALL and K. A. ALMALKI: Geochemistry of Late Cambrian-Early Ordovician fluvial to shallow marine sandstones, western Tasmania, Australia: Implications for provenance, weathering, tec- tonic settings, and chemostratigraphy | 197 |
| <i>EXPRESS LETTER</i> | |
| H. KATO and T. YAMADA: Controlling factors in stalagmite oxygen isotopic composition and the paleoprecipitation record for the last 1,100 years in Northeast Japan | e1 |